

BINARY BLACK HOLE MERGER WAVEFORMS IN THE EXTREME MASS RATIO LIMIT

THIBAUT DAMOUR

Institut des Hautes Etudes Scientifiques, 35 route de Chartres, 91440 Bures-sur-Yvette, France

ALESSANDRO NAGAR

Dipartimento di Fisica, Politecnico di Torino, and INFN, sezione di Torino, Torino, Italy

We discuss the transition from quasi-circular inspiral to plunge of a system of two non-rotating black holes of masses m_1 and m_2 in the extreme mass ratio limit $m_1 m_2 \ll (m_1 + m_2)^2$. In this limit, we compare the merger waveforms obtained by two different methods: a *numerical* (Regge-Wheeler-Zerilli) one, and an *analytical* (Effective One Body) one. This is viewed as a contribution to the matching between analytical and numerical methods.

1. Introduction

The last months have witnessed a decisive advance in Numerical Relativity, with different groups being able to simulate the merger of two black holes of comparable masses.¹ Since such binary black holes systems (of a total mass $\sim 30M_\odot$) are believed to be among the most promising sources of gravitational waves for the ground based detectors like LIGO and VIRGO, this breakthrough raises the hope to have, for the first time, a reliable estimate of the complete waveform by joining together Post-Newtonian (PN) and Numerical Relativity results. We recall that PN techniques have provided us with high-order results for describing the motion and radiation² of binary systems, and that further techniques have been proposed for *resumming* the PN results,^{3,4} thereby allowing an analytical description of the gravitational waveform emitted during the transition between inspiral and plunge, and even during the subsequent merger and ringdown phases. We now face the important task of constructing accurate *complete waveforms* by *matching* together the information contained in *Post-Newtonian* and *Numerical Relativity* results. We view the present work as a contribution towards this goal (see also Refs.^{5,6}).

The present work belongs to a scientific lineage which was started by Regge and Wheeler,⁷ Zerilli,⁸ Davis, Ruffini, Press and Price⁹ and Davis, Ruffini and Tiomno.¹⁰ Refs.^{9,10} studied the gravitational wave emission due to the *radial plunge* (from infinity) of a particle into a Schwarzschild black hole, as a model for the head-on collision of two black holes in the extreme mass ratio limit. Here we consider, for the first time, the *transition* from the quasi-circular adiabatic inspiral phase to the plunge phase in extreme-mass-ratio binary black hole systems. We shall be able to do that by getting round a limitation of the original Regge-Wheeler-Zerilli test-particle approach: that of requiring the test particle to follow an exact *geodesic* of the Schwarzschild background. We bypass this stumbling block by using an improved form of PN theory: the Effective One Body (EOB) approach to the general relativistic two body dynamics. This approach has been recently proposed to study




Fig. 1. *Left panel:* Gravitational waveforms (real and imaginary part) for $l = m = 2$ generated by a plunge from an initial separation $r = 7M$. *Right panel:* Comparison between the corresponding gravitational-wave phase and that obtained from analytically matching a 3PN improved quadrupole-type formula to a superposition of quasi-normal modes.

the transition from inspiral to plunge in the *comparable-mass* case.⁴ It describes the dynamics of a binary system in terms of two separate ingredients: (i) a Hamiltonian $H_{\text{EOB}}(M, \mu)$ describing the *conservative* part of the relative dynamics, and (ii) a non-Hamiltonian supplementary force $\mathcal{F}_{\text{EOB}}(M, \mu)$ approximately describing the reaction to the loss of energy and angular momentum along quasi-circular orbits. [Here, $M \equiv m_1 + m_2$, and $\mu = m_1 m_2 / M$.] The badly convergent PN-Taylor series giving the angular momentum flux is *resummed* by means of Padé approximants.³

2. Results

We summarize here our tools and main results (see Refs.^{12,13} for more information). We describe, in the extreme mass ratio limit, the relative dynamics of the binary system of two non-rotating black holes by that of a “particle” (of mass $\mu \ll M$) that follows a non-geodesic inspiral driven by radiation-reaction (until a quasi-geodesic plunge) on a (quasi-)Schwarzschild background of mass M . We then use the multipolar Regge-Wheeler-Zerilli perturbation theory around such a black hole to compute the gravitational wave emission in the approximation $\mu \ll M$. Einstein equations then lead to two (even/odd) *decoupled* wave-like equations (with source terms $S_{\ell m}^{(e/o)} \propto \mu$) for two (even/odd) master functions $\Psi_{\ell m}^{(e/o)}$.¹¹

Fig. 1 shows the $l = m = 2$ waveform generated by a binary system with initial relative separation $r = 7M$ ^a. After a “chirp-like” increase of the frequency and modulus during the inspiral, the latter reaches a maximum when the particle crosses the light ring ($r = 3M$) and eventually the waveform decays in a quasi-normal mode ringing phase.

^aNote that the other multipoles give a non-negligible contribution, due to the strong asymmetry of the system.

The so-constructed “exact” *numerical* waveforms are then compared with semi-*analytical* ones, constructed within the EOB framework and philosophy. We recall that the basic idea of the EOB framework is to produce quasi-analytical waveforms by patching together a quadrupole-type waveform during the inspiral and plunge to a QNM-type waveform after merger. We give an example of this numerical-analytical comparison in the right panel of Fig. 1. The gravitational wave phases ϕ_{gw}^{22} obtained by two different methods, one numerical and the other semi-analytical, are compared. In the first method, ϕ_{gw}^{22} is given from time-integration of the instantaneous gravitational wave frequency obtained as $M\omega_{\text{gw}}^{22} = -\Im(\dot{\Psi}_{22}^{(e)}/\Psi_{22}^{(e)})$. The second method computes an approximate waveform in the following way: before crossing the light ring one uses a (Padé resummed) 3PN-improved quadrupole-type formula to compute the waveform from the EOB dynamics. After crossing the light ring, the previous quadrupole-type signal, taken in the quasi-circular (QC) approximation, is *matched* to a superposition of the first five QNMs of the black hole. Then one computes the phase of the matched analytical waveform by integrating the corresponding instantaneous gravitational wave frequency. The maximum difference between the “exact” phase and the “effective one body” phase turns out to be less than 3% of a cycle (see inset).

References

1. F. Pretorius, *Phys. Rev. Lett.* **95**, 121101 (2005); M. Campanelli, C.O. Lousto, P. Marronetti, and Y. Zlochower, *Phys. Rev. Lett.* **96** 111101 (2006); J.G. Baker, J. Centrella, D.I. Choi, M. Koppitz, and J. van Meter *Phys. Rev. Lett.* **96** 111102 (2006); M. Campanelli, C.O. Lousto, and Y. Zlochower, *Phys. Rev. D* **74**, 084023 (2006); J. Gonzalez, U. Sperhake, B. Bruegmann, M. Hannam, S. Husa, *arXiv:gr-qc/0610154*.
2. T. Damour, P. Jaranowski and G. Schäfer, *Phys. Lett. B* **513**, 147 (2001); L. Blanchet, T. Damour, and G. Esposito-Farèse, *Phys. Rev. D* **69**, 124007 (2004); L. Blanchet, T. Damour, G. Esposito-Farèse and B.R. Iyer, *Phys. Rev. D* **71**, 124004 (2005); L. Blanchet, *Living Rev. Relativity* **9**, 4. URL: <http://www.livingreviews.org/lrr-2006-4> (2006).
3. T. Damour, B.R. Iyer and B.S. Sathyaprakash, *Phys. Rev. D*, **57**, 885 (1998).
4. A. Buonanno and T. Damour, *Phys. Rev. D* **59**, 084006 (1999); *ibid.*, **62**, 064015 (2000).
5. A. Buonanno, G.B. Cook, and F. Pretorius, *arXiv: gr-qc/0610122* (2006).
6. J. G. Baker, J. R. van Meter, S. T. McWilliams, J. Centrella and B. J. Kelly, *arXiv:gr-qc/0612024*.
7. T. Regge and J.A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
8. F.J. Zerilli, *Phys. Rev. Lett.* **24**, 737 (1970).
9. M. Davis, R. Ruffini, W.H. Press, and R.H. Price, *Phys. Rev. Lett.* **27**, 1466 (1971).
10. M. Davis, R. Ruffini, and J. Tiomno, *Phys. Rev. D* **5**, 2932 (1972).
11. A. Nagar and L. Rezzolla, *Class. Q. Grav.* **22**, R167 (2005); erratum, **23**, 4297 (2006).
12. A. Nagar, T. Damour and A. Tartaglia, in *Class. Q. Grav.* special issue based on *New Frontiers in Numerical Relativity* conference, Golm (Germany), July 17-22 2006, *arXiv: gr-qc/0612096*.
13. T. Damour and A. Nagar, *in preparation*.